

Well Child and Immunization Schedule (0-5 Years)

Initial Visit:	3-10 days after birth; No immunizations.
One month:	No immunizations.
Two months:	# 1 Pediarix (DTaP, IPV, and Hepatitis B) # 1 Hib # 1 Pevnar # 1 Rotateq
Four months:	# 2 Pediarix # 2 Hib # 2 Pevnar # 2 Rotateq
Six months:	# 3 Pediarix # 3 Hib # 3 Pevnar # 3 Rotateq
Nine months:	Catch-up immunizations if needed Hemoglobin (fingerstick)
Twelve months:	# 1 MMR (per provider) # 1 Varicella # 4 Pevnar (per provider) # 1 Hep A (per provider)
Fifteen months:	# 1 MMR (if not already done) # 4 Hib # 4 Pevnar (if not already done) DTaP
Eighteen months:	Catch-up on any missing/late immunizations # 2 Hep A (per provider)
Two years:	Catch-up on any missing/late immunizations

Influenza (flu) vaccine recommended for children 6 months and up.

Annual physicals with booster immunizations at school age (4-6yr) **MMR, DTaP, IPV,** and **Varicella**

Annual physicals with booster immunizations at middle school (11-12yr) **Tdap** and **Menactra**

DTaP – Diphtheria, Tetanus, Acellular Pertussis

Hib – Haemophilus Influenza Type B

IPV – Polio

Rotateq – Rotavirus

Pevnar – Pneumococcal Conjugate

MMR – Measles, Mumps, and Rubella

Varicella – Chicken Pox

Tdap – Tetanus, Diphtheria, Acellular Pertussis

Menactra - Meningococcal disease